arXiv:math/0506476v6 [math.DS] 22 Aug 2005

On coding with Feller contractive Markov systems

Ivan Werner
Email: ivan_werner@pochta.ru

February 1, 2008

Abstract

We continue development of the theory of contractive Markov systems initiated in [15]. In this paper, we construct the coding map for Feller contractive Markov systems. This allows us to prove a generalization of Ledrappier's Theorem [10] and to show the existence of observable invariant measures for Feller contractive Markov system.

MSC: 60J05, 37D35, 37A50, 37H99, 28A80
Keywords: contractive Markov systems (CMS), iterated function systems with place-dependent probabilities, random systems with complete connections, g-measures, Markov chains, equilibrium states, fractals.

1 Introduction

In [15], the author introduced a concept of contractive Markov systems (CMSs). It unifies some structures which arise in different areas of mathematics, such as *finite Markov chains*; *learning models* [6]; *g-measures* [8], [10], [13], [7]; *iterated function systems with place-dependent probabilities* [1], [3]; *graph-directed constructions of 'fractal' sets* [11]. It can be seen as a part of a general theory of *random systems with complete connections* [5]. However, the proposed structure seems already to have a sufficient level of generality which allows one to see some "forest" behind the "trees" and to contribute some new results also to other areas of mathematics such as *ergodic theory* [14], [9].

So, what is a Markov system. It is a structure on a metric space which generates a discrete-time Markov process on it.

Let $K_1, K_2, ..., K_N$ be a partition of a metric space K into non-empty Borel subsets (we do not exclude the case $N = 1$). Furthermore, for each $i \in \{1, 2, ..., N\}$, let

$$w_{i1}, w_{i2}, ..., w_{iL_i} : K_i \longrightarrow K$$

be a family of Borel measurable maps such that for each $j \in \{1, 2, ..., L_i\}$ there exists $n \in \{1, 2, ..., N\}$ such that $w_{ij}(K_i) \subset K_n$ (Fig. 1). Finally, for each $i \in \{1, 2, ..., N\}$, let

$$p_{i1}, p_{i2}, ..., p_{iL_i} : K_i \longrightarrow \mathbb{R}^+$$

be a family of positive Borel measurable probability functions (associated with the maps), i.e. $p_{ij} > 0$ for all j and $\sum_{j=1}^{L_i} p_{ij}(x) = 1$ for all $x \in K_i$.



Fig. 1. A Markov system.

Definition 1 We call $V := \{1, ..., N\}$ *the set of vertices* and the subsets $K_1, ..., K_N$ are called *the vertex sets*. Further, we call

$$E := \{(i, n_i) : i \in \{1, ..., N\}, n_i \in \{1, ..., L_i\}\}$$

the set of edges and we use the following notations:

$$p_e := p_{in} \text{ and } w_e := w_{in} \text{ for } e := (i, n) \in E.$$

Each edge is provided with a direction (an arrow) by marking *an initial vertex* through the map

$$\begin{array}{rcl} i : E & \longrightarrow & V \\ (j, n) & \longmapsto & j. \end{array}$$

The terminal vertex $t(j, n) \in V$ of an edge $(j, n) \in E$ is determined by the corresponding map through

$$t((j, n)) \quad := \quad k \; :\Longleftrightarrow \quad w_{jn}(K_j) \subset K_k.$$

We call the quadruple $G := (V, E, i, t)$ *a directed (multi)graph* or *digraph*. A sequence (finite or infinite) $(..., e_{-1}, e_0, e_1, ...)$ of edges which corresponds to a walk along the arrows of the digraph (i.e. $t(e_k) = i(e_{k+1})$) is called *a path*.

Definition 2 We call the family $\mathcal{M} := \left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ a *Markov system.*

Definition 3 A Markov system is called *irreducible* or *aperiodic* if and only if its directed graph is irreducible or aperiodic respectively.

Definition 4 (CMS) We call a Markov system $\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ *contractive* with an *average contracting rate* $0 < a < 1$ iff it satisfies the following *condition of contractiveness on average*:

$$\sum_{e\in E} p_e(x)d(w_e(x), w_e(y)) \leq ad(x,y) \text{ for all } x,y \in K_i \text{ and } i \in \{1,...,N\} \tag{1}$$

(it is understood here that p_e's are extended on the whole space by zero and w_e's arbitrarily). This condition was discovered by Richard Isaac in 1962 for the case $N = 1$ [6].

Remark 1 Note that also the situation applies where each vertex set K_i has its own metric d_i. In this case, one can set

$$d(x,y) = \begin{cases} d_i(x,y) & \text{if } x,y \in K_i \\ \infty & \text{otherwise} \end{cases}$$

and use the convention $0 \times \infty = 0$.

A Markov system $\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ determines a Markov operator U on the set of all bounded Borel measurable functions $\mathcal{L}^0(K)$ by

$$Uf := \sum_{e\in E} p_e f \circ w_e \text{ for all } f \in \mathcal{L}^0(K)$$

and its adjoint operator U^* on the set of all Borel probability measures $P(K)$ by

$$U^*\nu(f) := \int U(f)d\nu \text{ for all } f \in \mathcal{L}^0(K) \text{ and } \nu \in P(K).$$

We say a probability measure μ is an *invariant probability measure* of the Markov system if and only if it is a stationary initial distribution of the associated Markov process, i.e.

$$U^*\mu = \mu.$$

A Borel probability measure μ is called *attractive* measure of the CMS if and only if

$${U^*}^n\nu \overset{w^*}{\to} \mu \text{ for all } \nu \in P(K),$$

where w^* means weak* convergence. Note that an attractive probability measure is a unique invariant probability measure of the CMS if U maps continuous functions on continuous functions.

We shall denote the Borel σ-algebra on a set X by $\mathcal{B}(X)$.

In [16], we constructed an important tool for CMSs called the *coding map*. We showed that it is defined almost everywhere with respect to a natural measure on the *code space* if any two probability measures representing Markov processes with Dirac initial distributions concentrated on some points in the same vertex set are absolutely continuous. In [18], we showed that this condition also implies the uniqueness of the invariant probability measure. Moreover, we showed that this condition is satisfied if the probability functions $p_e|_{K_{i(e)}}$ have square summable variation, i.e. for any $b > 0$ and $0 < c < 1$,

$$\sum_{n=0}^{\infty} \phi^2\left(bc^n\right) < \infty,$$

where ϕ is *the modulus of uniform continuity* of the probability function, i.e.

$$\phi(t) := \sup\{|p_e(x) - p_e(y)| : d(x,y) \leq t,\ x,y \in K_{i(e)}\}.$$

The condition of the square summability of variation was introduced by A. Johansson and A. Öberg [7].

In this paper, we discuss contractive Markov systems which only have the Feller property, i.e. the Markov operator U maps the set of continuous functions into itself. For that to be satisfied, one needs only the continuity of the probability functions.

Example 1 Consider two maps $w_0(x) := 9/10x$ and $w_1(x) := -x + 1/e$ for all $x \in [0, 1/e]$ with probability functions given as follows. Let $\alpha, \delta > 0$ such that $\alpha + \delta < 1$. Set

$$p_0(x) = \begin{cases} \frac{\alpha}{\sqrt{\log\frac{1}{x}}} + \delta & \text{if } 0 < x \leq \frac{1}{e} \\ \delta & \text{if } x = 0. \end{cases}$$

and $p_1(x) := 1 - p_0(x)$ for all $x \in [0, 1/e]$. Then a simple calculation shows that $([0, 1/e], w_i, p_i)_{i\in\{0,1\}}$ defines a CMS with an average contraction rate $1 - 9/10\delta$. Obviously, this CMS has the Feller property. Let ϕ be the modulus of uniform continuity of p_0 and $0 < c < 1/e$. Then

$$\phi(c^n) = \frac{\alpha}{\sqrt{n\log\frac{1}{c}}} \text{ for all } n \in \mathbb{N}.$$

Hence p_0 does not have a square summable variation. Therefore, we do not know whether this CMS has a unique invariant Borel probability measure. It was shown by Berger, Hoffman and Sidoravicius [2] that the condition of the square summability of variation is tight, in the sense that for any $\epsilon > 0$ there exists a g-function with a summable variation to the power $2 + \epsilon$ such that the corresponding CMS on a one-sided full shift has several invariant Borel probability measures.

In this paper, we intend to show that the coding map still can be defined if the probabilities are merely uniformly continuous (Proposition 2). This opens a way of extending the thermodynamic formalism proposed in [17], and further developed and successfully used in [18], on the class of Feller CMSs. This leads to the generalization of Ledrappier's Theorem [10] (Theorem 2) and allows us to prove the existence of observable invariant measures for Feller CMSs (Corollary 1).

2 Main part

Let $\mathcal{M} := \left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ be a contractive Markov system with an average contracting rate $0 < a < 1$. We assume that: (K, d) is a complete separable metric space; the family $K_1, ..., K_N$ partitions K into non-empty open subsets; each probability function $p_e|_{K_{i(e)}}$ is uniformly continuous and bounded away from zero by $\delta > 0$; the set of edges E is finite and the map $i : E \longrightarrow V$ is surjective.

Note that the assumptions imply that the CMS has the Feller property.

Let $P(\mathcal{M})$ be the set of all invariant Borel probability measures of $\mathcal{M}$.

Proposition 1 *(i)* $P(\mathcal{M}) \neq \emptyset$.
(ii) For every $\mu \in P(\mathcal{M})$,

$$\sum_{i=1}^{N}\int_{K_i} d(x, x_i)\ d\mu(x) < \infty \textit{ for all } x_i \in K_i,\ i = 1, ..., N. \tag{2}$$

Proof. See [4]. □

Note that if (2) holds for some $x_j \in K_j$, $j = 1, ..., N$, by the triangle inequality, it holds for all $x_j \in K_j$, $j = 1, ..., N$.

Now, we are going to construct the coding map with respect to the generalized Markov measures associated with invariant measures of $\mathcal{M}$.

2.1 Coding map

Let $\Sigma := \{(..., \sigma_{-1}, \sigma_0, \sigma_1, ...) : e_i \in E\ \forall i \in \mathbb{Z}\}$. We shall denote by d' the metric on Σ defined by $d'(\sigma, \sigma') := (1/2)^k$ where k is the largest integer with $\sigma_i = \sigma'_i$ for all $|i| < k$. Denote by S the left shift map on Σ. We call the set ${}_m[e_m, ..., e_n] := \{\sigma \in \Sigma :\ \sigma_i = e_i \text{ for all } m \leq i \leq n\}$, for $m \leq n \in \mathbb{Z}$, a *cylinder set*. Let μ be an invariant Borel probability measure of $\mathcal{M}$. We call the shift invariant Borel probability measure M on Σ given by

$$M\left({}_m[e_1, ..., e_k]\right) := \int p_{e_1}(x)p_{e_2}(w_{e_1}x)...p_{e_k}(w_{e_{k-1}} \circ ... \circ w_{e_1}x)d\mu(x),$$

for every cylinder set ${}_m[e_1, ..., e_k]$, the *generalized Markov measure* (see [16]).

Proposition 2 *Let $\mu \in P(\mathcal{M})$. Let M be the generalized Markov measure associated with $\mathcal{M}$ and μ. Let $y_i,\ z_i \in K_i$ for all $i = 1, ..., N$. Then the following hold.*
(i)

$$\lim_{m\to-\infty} d\left(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(z_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(y_{i(\sigma_m)})\right) = 0\ M\text{-}a.e..$$

(ii)

$$F_{z_1...z_N} := \lim_{m\to-\infty} w_{\sigma_0}\circ ...\circ w_{\sigma_m}(z_{i(\sigma_m)})\ \text{exists}\ M\text{-}a.e.,$$

and by (i) $F_{z_1...z_N} = F_{y_1...y_N}$ M*-a.e..*
(iii) There exists a sequence of closed subsets $Q_1 \subset Q_2 \subset ... \subset \Sigma$ *with* $\sum_{k=1}^{\infty} M(\Sigma\setminus Q_k) < \infty$ *such that* $F_{z_1...z_N}|_{Q_k}$ *is locally Hölder-continuous with the same Hölder-constants for all* $k \in \mathbb{N}$*, i.e. there exist* $\alpha, C > 0$ *such that for every* k *there exists* $\delta_k > 0$ *such that*

$$\sigma, \sigma' \in Q_k\ \text{with}\ d'(\sigma,\sigma') \le \delta_k\ \Rightarrow\ d(F_{z_1...z_N}(\sigma), F_{z_1...z_N}(\sigma')) \le C d'(\sigma,\sigma')^{\alpha}.$$

Proof. Fix $x_j \in K_j$ for all $j = 1, ..., N$.

(i) First observe that, for all $m \le 0$,

$$\begin{aligned}
& \int d(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(y_{i(\sigma_m)}))\ dM \\
= & \sum_{e_m,...,e_0} M({}_m[e_m,...,e_0]) d(w_{e_0}\circ ...\circ w_{e_m}(x_{i(e_m)}), w_{e_0}\circ ...\circ w_{e_m}(y_{i(e_m)})) \\
= & \int \sum_{e_m,...,e_0} p_{e_m}(x)...p_{e_0}(w_{e_{-1}}\circ ...\circ w_{e_m}(x)) \\
& \times d(w_{e_0}\circ ...\circ w_{e_m}(x_{i(e_m)}), w_{e_0}\circ ...\circ w_{e_m}(y_{i(e_m)}))\ d\mu(x) \\
\le & \int \sum_{e_m,...,e_0} p_{e_m}(x)...p_{e_0}(w_{e_{-1}}\circ ...\circ w_{e_m}(x)) \\
& \times d(w_{e_0}\circ ...\circ w_{e_m}(x_{i(e_m)}), w_{e_0}\circ ...\circ w_{e_m}(x))\ d\mu(x) \\
& + \int \sum_{e_m,...,e_0} p_{e_m}(x)...p_{e_0}(w_{e_{-1}}\circ ...\circ w_{e_m}(x)) \\
& \times d(w_{e_0}\circ ...\circ w_{e_m}(x), w_{e_0}\circ ...\circ w_{e_m}(y_{i(e_m)}))\ d\mu(x)
\end{aligned}$$

$$
\begin{aligned}
\leq\ & \sum_{j=1}^{N}\int_{K_j}\sum_{e_m,\dots,e_0} p_{e_m}(x)...p_{e_0}(w_{e_{-1}}\circ ...\circ w_{e_m}(x)) \\
& \times d(w_{e_0}\circ ...\circ w_{e_m}(x_j), w_{e_0}\circ ...\circ w_{e_m}(x))\ d\mu(x) \\
& + \sum_{j=1}^{N}\int_{K_j}\sum_{e_m,\dots,e_0} p_{e_m}(x)...p_{e_0}(w_{e_{-1}}\circ ...\circ w_{e_m}(x)) \\
& \times d(w_{e_0}\circ ...\circ w_{e_m}(x), w_{e_0}\circ ...\circ w_{e_m}(y_j))\ d\mu(x) \\
\leq\ & a^{-m+1}\sum_{j=1}^{N}\int_{K_j} d(x_j,x)\ d\mu(x) + a^{-m+1}\sum_{j=1}^{N}\int_{K_j} d(x,y_j)\ d\mu(x).
\end{aligned}
$$

Hence, there exists $C>0$ such that

$$\int d(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(y_{i(\sigma_m)}))\ dM \leq a^{-m+1}C \text{ for all } m\leq 0.$$

Set

$$A_m := \left\{\sigma\in\Sigma:\ d\left(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(y_{i(\sigma_m)})\right) > a^{\frac{-m+1}{2}}C\right\}$$

for all $m\leq 0$. Then, by the Markov inequality,

$$M(A_m)\leq a^{\frac{-m+1}{2}} \text{ for all } m\leq 0.$$

Now, let

$$A := \bigcap_{l=0}^{-\infty}\bigcup_{m=l}^{-\infty} A_m.$$

Then

$$M(A)\leq \sum_{m\leq l} a^{\frac{-m+1}{2}} \text{ for all } l\leq 0.$$

Hence, $M(A)=0$, but for all $\sigma\in\Sigma\setminus A$,

$$\lim_{m\to-\infty} d\left(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(y_{i(\sigma_m)})\right) = 0. \tag{3}$$

Analogously, holds

$$\lim_{m\to-\infty} d\left(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(x_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(z_{i(\sigma_m)})\right) = 0\ M\text{-a.e..}$$

Hence, by the triangle inequality,

$$\lim_{m\to-\infty} d\left(w_{\sigma_0}\circ ...\circ w_{\sigma_m}(y_{i(\sigma_m)}), w_{\sigma_0}\circ ...\circ w_{\sigma_m}(z_{i(\sigma_m)})\right) = 0\ M\text{-a.e..}$$

(ii) Let us abbreviate $Y_m(\sigma) := w_{\sigma_0} \circ ... \circ w_{\sigma_m}(x_{i(\sigma_m)})$ for $\sigma \in \Sigma$ and $m \leq 0$. Then similarly as in the proof of (i), for all $m \leq 0$,

$$\begin{aligned}
& \int d(Y_m, Y_{m-1})\ dM \\
= & \int \sum_{e_{m-1},...,e_0} p_{e_{m-1}}(x)...p_{e_0}(w_{e_{-1}} \circ ... \circ w_{e_{m-1}}(x)) \\
& \times d(w_{e_0} \circ ... \circ w_{e_m}(x_{i(e_m)}), w_{e_0} \circ ... \circ w_{e_{m-1}}(x_{i(e_{m-1})}))\ d\mu(x) \\
\leq & \int \sum_{e_{m-1},...,e_0} p_{e_{m-1}}(x)...p_{e_0}(w_{e_{-1}} \circ ... \circ w_{e_{m-1}}(x)) \\
& \times d(w_{e_0} \circ ... \circ w_{e_m}(x_{i(e_m)}), w_{e_0} \circ ... \circ w_{e_{m-1}}(x))\ d\mu(x) \\
& + \int \sum_{e_{m-1},...,e_0} p_{e_{m-1}}(x)...p_{e_0}(w_{e_{-1}} \circ ... \circ w_{e_{m-1}}(x)) \\
& \times d(w_{e_0} \circ ... \circ w_{e_{m-1}}(x), w_{e_0} \circ ... \circ w_{e_{m-1}}(x_{i(e_{m-1})}))\ d\mu(x) \\
\leq & \sum_{j=1}^{N} \int \sum_{(e_{m-1},...,e_0),\ t(e_{m-1})=j=i(e_m)} p_{e_{m-1}}(x)...p_{e_0}(w_{e_{-1}} \circ ... \circ w_{e_{m-1}}(x)) \\
& \times d(w_{e_0} \circ ... \circ w_{e_m}(x_j), w_{e_0} \circ ... \circ w_{e_{m-1}}(x))\ d\mu(x) \\
& + \sum_{j=1}^{N} \int_{K_j} \sum_{e_{m-1},...,e_0} p_{e_{m-1}}(x)...p_{e_0}(w_{e_{-1}} \circ ... \circ w_{e_{m-1}}(x)) \\
& \times d(w_{e_0} \circ ... \circ w_{e_{m-1}}(x), w_{e_0} \circ ... \circ w_{e_{m-1}}(x_j))\ d\mu(x) \\
\leq & \ a^{-m+1} \sum_{j=1}^{N} \int \sum_{e_{m-1}, t(e_{m-1})=j} p_{e_{m-1}}(x) d(x_j, w_{e_{m-1}}(x))\ d\mu(x) \\
& + a^{-m+2} \sum_{j=1}^{N} \int_{K_j} d(x, x_j)\ d\mu(x) \\
= & \ a^{-m+1} \int U\left(\sum_{j=1}^{N} 1_{K_j} d(x_j, .)\right)(x)\ d\mu(x) + a^{-m+2} \sum_{j=1}^{N} \int_{K_j} d(x, x_j)\ d\mu(x) \\
= & \ a^{-m+1} \int \sum_{j=1}^{N} 1_{K_j} d(x_j, x)\ d\mu(x) + a^{-m+2} \sum_{j=1}^{N} \int_{K_j} d(x, x_j)\ d\mu(x)
\end{aligned}$$

Hence, there exists $C > 0$ such that,

$$\int d(Y_m, Y_{m-1})\ dM \leq a^{-m+1} C \text{ for all } m \leq 0.$$

Let

$$B_{m-1} := \left\{ \sigma \in \Sigma :\ d(Y_m, Y_{m-1}) > a^{\frac{-m+1}{2}} C \right\} \text{ for all } m \leq 1.$$

Then, by the Markov inequality,

$$M(B_{m-1}) \leq a^{\frac{-m+1}{2}} \text{ for all } m-1 \leq 0.$$

Let

$$B := \bigcap_{l=0}^{-\infty} \bigcup_{m=l}^{-\infty} B_m.$$

Then

$$M(B) \leq \sum_{m \leq l} a^{\frac{-m}{2}} \text{ for all } l \leq 0. \tag{4}$$

Hence, $M(B) = 0$, and for all $\sigma \in \Sigma \setminus B$, the sequence $(Y_m(\sigma))_{m\leq 0}$ is Cauchy. Thus

$$F_{x_1 \ldots x_N} := \lim_{m \to -\infty} w_{\sigma_0} \circ \ldots \circ w_{\sigma_m}(x_{i(\sigma_m)}) \text{ exists } M\text{-a.e.},$$

and by (3) $F_{y_1 \ldots y_N}$ also exists M-a.e..

(iii) Let

$$Q_l := \Sigma \setminus \bigcup_{m=l}^{-\infty} B_m \text{ for all } l \in \mathbb{Z} \setminus \mathbb{N},$$

where B_m as in proof of part (ii). Then $Q_l \subset Q_{l-1}$ for all $l \in \mathbb{Z} \setminus \mathbb{N}$. Since every Y_m is continuous, B_m is open for all $m \in \mathbb{Z} \setminus \mathbb{N}$, and so, every Q_l is closed. Since $M(B_m) \leq a^{\frac{-m}{2}}$ for all $m \leq 0$,

$$\sum_{l=0}^{-\infty} M(\Sigma \setminus Q_l) = \sum_{l=0}^{-\infty} \sum_{m=l}^{-\infty} a^{\frac{-m}{2}} < \infty.$$

Now, let $\sigma, \sigma' \in Q_l$ for some l. Then

$$d\left(Y_m(\sigma), Y_{m-1}(\sigma)\right) \leq a^{\frac{-m+1}{2}} C \text{ for all } m-1 \leq l.$$

Therefore, by the triangle inequality,

$$d\left(Y_m(\sigma), Y_{m-k}(\sigma)\right) \leq \sum_{i=m}^{-\infty} a^{\frac{-i+1}{2}} C = C \frac{1}{1-\sqrt{a}}\; a^{\frac{|m-1|}{2}} \text{ for all } m-1 \leq l,\ k \geq 1.$$

Hence

$$d\left(Y_m(\sigma), F_{x_1 \ldots x_N}(\sigma)\right) \leq C \frac{1}{1-\sqrt{a}}\; a^{\frac{|m-1|}{2}} \text{ for all } m-1 \leq l.$$

Analogously,

$$d\left(Y_m(\sigma'), F_{x_1 \ldots x_N}(\sigma')\right) \leq C \frac{1}{1-\sqrt{a}}\; a^{\frac{|m-1|}{2}} \text{ for all } m-1 \leq l.$$

Now, let $\delta_l = (1/2)^{|l-1|}$, $\alpha := \log\sqrt{a}/\log(1/2)$ and $d'(\sigma,\sigma') = (1/2)^{|m-1|}$ for some $m-1 \leq l$. Then $Y_m(\sigma') = Y_m(\sigma)$. Therefore,

$$d\left(F_{x_1\ldots x_N}(\sigma), F_{x_1\ldots x_N}(\sigma')\right) \leq \frac{2C}{1-\sqrt{a}}\, a^{\frac{|m-1|}{2}} = \frac{2C}{1-\sqrt{a}}\, d'(\sigma,\sigma')^{\alpha}.$$

□

Remark 2 Note that the Feller property of $\mathcal{M}$ was *not* used in the proof of Proposition 2. It only insures that $P(\mathcal{M})$ is not empty.

Definition 5 We call the map

$$F_{x_1\ldots x_N} : \Sigma \longrightarrow K,$$

which is well defined by Proposition 2, the *coding map* of the CMS.

Remark 3 Proposition 2 generalizes Corollary 1 from [16].

2.2 Thermodynamic formalism

In this subsection, we improve some results from [17] and [18]. In [17], we initiated the development of the thermodynamic formalism for contractive Markov systems. It was shown that it goes beyond the scope of the existing theory of equilibrium states. Namely, in case of contractive Markov systems we have to consider local energy functions which are not upper semicontinuous. In [18], we successfully used the developed thermodynamic formalism for the proof of the uniqueness of the invariant probability measure for CMSs satisfying a 'regularity condition' which holds if the probability functions $p_e|_{K_{i(e)}}$ have square summable variation.

Definition 6 Let X be a metric space and T a continuous transformation on it. Denote by $P(X)$ the set of all Borel probability measures on X and by $P_T(X)$ the set of all T-invariant Borel probability measures on X. We call a Borel measurable function $f : X \longrightarrow [-\infty, 0]$ an *energy function.* Suppose that T has a finite topological entropy, i.e. $\sup_{\Theta\in P_T(X)} h_\Theta(T) < \infty$, where $h_\Theta(T)$ is the Kolmogorov-Sinai entropy of T with respect to measure Θ. We call

$$P(f) = \sup_{\Theta\in P_T(X)} \left(h_\Theta(T) + \Theta(f)\right)$$

the *pressure* of f. We call $\Lambda \in P_T(X)$ *an equilibrium state* for f iff

$$h_\Lambda(T) + \Lambda(f) = P(f).$$

Let's denote the set of all equilibrium states for f by $ES(f)$.

Now, we define the energy function for the CMS $\mathcal{M}$.

Definition 7 Fix $x_i \in K_i$ for each $i = 1, ..., N$. In what follows, let's denote $F_{x_1 \dots x_N}$ simply by F. Let

$$\Sigma_G := \{\sigma \in \Sigma : \ t(\sigma_j) = i(\sigma_{j+1}) \ \forall j \in \mathbb{Z}\},$$

$$D := \{\sigma \in \Sigma_G : \ \lim_{m \to -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m} x_{i(\sigma_m)} \text{ exists}\}$$

and

$$Y := \bigcap_{i=-\infty}^{\infty} S^i(D).$$

Set

$$u(\sigma) = \begin{cases} \log p_{\sigma_1} \circ F(\sigma) & \text{if } \sigma \in Y \\ -\infty & \text{if } \sigma \in \Sigma \setminus Y. \end{cases} \tag{5}$$

Lemma 1 *(i) $F(\sigma)$ is defined for all $\sigma \in Y$ and Y is a shift invariant subset of Σ_G.*
(ii) For every $\mu \in P(\mathcal{M})$, $M(Y) = 1$, where M is the generalized Markov measure associated with the CMS and μ.
(iii) Suppose $\mathcal{M}$ is irreducible. Then Y is dense in Σ_G.

Proof. (i) is clear, by the definitions of F and Y. Since each map $w_e|_{K_{i(e)}}$ is continuous, $S(D) \subset D$. By Proposition 2, $M(D) = 1$. This implies (ii). If $\mathcal{M}$ is irreducible, then $\mu(K_i) > 0$ for all $i = 1, ..., N$ (e.g. Lemma 1 in [15]). Hence $M(O) > 0$ for every open $O \subset \Sigma_G$. This implies (iii). □

Remark 4 (i) Note that Y and F depend on the choice of x_i's. By Proposition 2 (i), Y changes only modulo M-zero set by a different choice of x_i's.
(ii) If all maps $w_e|_{K_{i(e)}}$ are contractive, then $Y = \Sigma_G$ and $F|_{\Sigma_G}$ is Hölder-continuous (easy to check).

Remark 5 As far as the author is aware, the rigorous mathematical theory of equilibrium states is developed only for upper semicontinuous energy functions f (see e.g. [9]). This condition insures that the convex space $ES(f)$ is non-empty and compact in the weak* topology. However, as the next example shows, u is not upper semicontinuous in general. Therefore, even the existence of an equilibrium state for u is not guaranteed by the existing theory of thermodynamic formalism.

Example 2 Let $(K, d) = (\mathbb{R}, |.|)$. Consider two maps

$$w_0(x) := \frac{1}{2}x, \ w_1(x) := 2x \text{ for all } x \in \mathbb{R}$$

with probability functions

$$p_0(x) := \frac{1}{6}\sin^2 x + \frac{17}{24}, \ p_1(x) := \frac{1}{6}\cos^2 x + \frac{1}{8} \text{ for all } x \in \mathbb{R}.$$

Then a simple calculation shows that $(\mathbb{R}, w_e, p_e)_{e\in\{0,1\}}$ defines a CMS with an average contracting rate 45/48. In this case, $\Sigma_G = \{0,1\}^{\mathbb{Z}}$. If we take $x = 0$ for the definition of Y, then, obviously, $Y = \Sigma_G$. Now, let $x \neq 0$. Let $N_{0n}(\sigma)$ and $N_{1n}(\sigma)$ be the numbers of zeros and ones in $(\sigma_{-n}, ..., \sigma_0)$ respectively for every $\sigma \in \Sigma_G$. Then, obviously, $\sigma \notin Y$ if $(N_{1n}(\sigma) - N_{0n}(\sigma)) \to \infty$. Hence, $Y \neq \Sigma_G$ and, by Lemma 1 (iii), Y is a dense shift invariant subset of Σ_G. Since Y is not closed, u is not upper semicontinuous.

Also, it is not difficult to see that in a general case there is no hope to find x_i such that u becomes upper semicontinuous, e.g. change w_1 to $w_1(x) = 2x + 1$, then, for any choice of x for the definition of Y, $Y \neq \Sigma_G$.

Before we move to the results on thermodynamic formalism for general CMSs, let us consider an important example, the thermodynamic formalism for which was elaborated by Ledrappier [10].

Example 3 Let $G := (V, E, i, t)$ be a finite irreducible directed (multi)graph. Let $\Sigma_G^- := \{(..., \sigma_{-1}, \sigma_0) : \ \sigma_m \in E \text{ and } t(\sigma_m) = i(\sigma_{m-1}) \ \forall m \in \mathbb{Z}\setminus\mathbb{N}\}$ (be *one-sided subshift of finite type* associated with G) endowed with the metric $d(\sigma, \sigma') := 2^k$ where k is the smallest integer with $\sigma_i = \sigma'_i$ for all $k < i \leq 0$. Let g be a positive, continuous function on Σ_G such that

$$\sum_{y\in T^{-1}(\{x\})} g(y) = 1 \text{ for all } x \in \Sigma_G$$

where T is the right shift map on Σ_G^-. Set $K_i := \left\{\sigma \in \Sigma_G^- : t(\sigma_0) = i\right\}$ for every $i \in V$ and, for every $e \in E$,

$$\tilde{w}_e(\sigma) := (..., \sigma_{-1}, \sigma_0, e),\ \tilde{p}_e(\sigma) := g(..., \sigma_{-1}, \sigma_0, e) \text{ for all } \sigma \in K_{i(e)}.$$

Obviously, maps $(\tilde{w}_e)_{e\in E}$ are contractions. Therefore, $\mathcal{M}_g := \left(K_{i(e)}, \tilde{w}_e, \tilde{p}_e\right)_{e\in E}$ defines a CMS. An invariant probability measure of such a CMS is called a *g-measure*. This notion was introduced by M. Keane [8]. Let U_g be the Markov operator associated with CMS $\mathcal{M}_g$. The following theorem characterizes the g-measures as equilibrium states for $\log g$.

Theorem 1 (Ledrappier, 1974) *Let $m \in P(\Sigma_G^-)$. Then the following are equivalent:*
(i) ${U_g}^* m = m$,
(ii) $m \in P_T(\Sigma_G^-)$ and $E_m(1_{{}_0[e]}|T^{-1}\mathcal{B}) = \tilde{p}_e \circ T$ m-a.e. for all $e \in E$ where $T^{-1}\mathcal{B}$ is the σ-algebra generated by cylinder sets of the form ${}_{-k}[e_{-k}, ..., e_{-1}] \subset \Sigma_G^-$,
(iii) $m \in P_T(\Sigma_G^-)$ and m is an equilibrium state for $\log g$ with $h_m(T) + m(\log g) = 0$.

Proof. See [10] or [13]. There, the g-measures are called the invariant measures of the *Ruelle operator* $\mathcal{L}_g$, given on the space of all Borel measurable functions

ϕ on Σ_G^- by

$$\mathcal{L}_g\phi(x) := \sum_{y\in T^{-1}(\{x\})} g(y)\phi(y) \text{ for all } x \in \Sigma_G^-.$$

However, $\mathcal{L}_g$ is nothing else but the Markov operator associated with CMS $\mathcal{M}_g$. In deed, for every function ϕ and $\sigma \in \Sigma_G^-$,

$$\begin{aligned}
\mathcal{L}_g\phi(\sigma) &= \sum_{e\in E:\ i(e)=t(\sigma_0)} g(...,\sigma_{-1},\sigma_0,e)\phi(...,\sigma_{-1},\sigma_0,e)\\
&= \sum_{e\in E} \tilde{p}_e(\sigma)\phi\circ\tilde{w}_e(\sigma)\\
&= U_g\phi(\sigma).
\end{aligned}$$

$\square$

Let $\pi : \Sigma_G \longrightarrow \Sigma_G^-$ be the natural projection. We call $\pi(\Lambda)$ *the natural restriction* of $\Lambda \in P(\Sigma_G)$ on Σ_G^-, and we call $\Lambda \in P(\Sigma_G)$ *the natural extension* of $\lambda \in P(\Sigma_G^-)$ if and only if $\pi(\Lambda) = \lambda$. Obviously, for every $\lambda \in P_T(\Sigma_G^-)$ the natural extension is unique. Observe that, in this example, the coding map $F = \pi$. The following proposition explains the relation between g-measures and the generalized Markov measures.

Proposition 3 *The generalized Markov measure associated with CMS $\mathcal{M}_g$ and a g-measure is the natural extension of the g-measure, or in other words, every g-measure is the image of the corresponding generalized Markov measure under the coding map.*

Proof. Let $m_g \in P(\Sigma_G^-)$ be a g-measure. Let $M_g \in P(\Sigma)$ be the generalized Markov measure associated with CMS $\mathcal{M}_g$ and m_g. Then, for every cylinder set ${}_{-k}[e_0,...,e_k] \subset \Sigma_G^-$,

$$\begin{aligned}
m_g({}_{-k}[e_0,...,e_k]) &= {U_g^*}^{k+1} m_g({}_{-k}[e_0,...,e_k])\\
&= \int \sum_{e\in E} \tilde{p}_e 1_{{}_{-k}[e_0,\dots,e_k]}\circ\tilde{w}_e d{U_g^*}^{k} m_g\\
&= \int \tilde{p}_{e_k} 1_{{}_{-k}[e_0,\dots,e_k]}\circ\tilde{w}_{e_k} d{U_g^*}^{k} m_g\\
&= \int \sum_{e\in E} \tilde{p}_e\tilde{p}_{e_k}\circ\tilde{w}_e 1_{{}_{-k}[e_0,\dots,e_k]}\circ\tilde{w}_{e_k}\circ\tilde{w}_e d{U_g^*}^{k-1} m_g\\
&= \int \tilde{p}_{e_{k-1}}\tilde{p}_{e_k}\circ\tilde{w}_{e_{k-1}} 1_{{}_{-k}[e_0,\dots,e_k]}\circ\tilde{w}_{e_k}\circ\tilde{w}_{e_{k-1}} d{U_g^*}^{k-1} m_g\\
&\quad .\\
&\quad .\\
&\quad .
\end{aligned}$$

$$\begin{aligned}
&= \int \tilde{p}_{e_0}\tilde{p}_{e_1}\circ\tilde{w}_{e_0}...\tilde{p}_{e_k}\circ\tilde{w}_{e_{k-1}}\circ...\circ\tilde{w}_{e_0} \\
&\quad \times 1_{{}_{-k}[e_0,...,e_k]}\circ\tilde{w}_{e_k}\circ\tilde{w}_{e_{k-1}}\circ...\circ\tilde{w}_{e_0}\,dm_g \\
&= \int \tilde{p}_{e_0}\tilde{p}_{e_1}\circ\tilde{w}_{e_0}...\tilde{p}_{e_k}\circ\tilde{w}_{e_{k-1}}\circ...\circ\tilde{w}_{e_0}\,dm_g \\
&= \pi\left(M_g\right)\left({}_{-k}[e_0,...,e_k]\right).
\end{aligned}$$

Hence, $m_g = \pi\left(M_g\right)$. □.

Now, we are going to prove a generalization of Theorem 1 which will relate invariant measures of CMS $\mathcal{M}$ with equilibrium states of u via the coding map.

Let $\mathcal{A}_0$ be the finite σ-algebra on Σ generated by the zero-time partition $\{{}_0[e] : e\in E\}$. Set $\mathcal{F} := \bigvee_{i=0}^{\infty} S^i\mathcal{A}_0$.

Theorem 2 *Let $\mu\in P(K)$. Then between the statements*
(i) $U^*\mu=\mu$,
(ii) $E_M(1_{{}_1[e]}|\mathcal{F}) = p_e\circ F$ M-a.e. for all $e\in E$, where M is the generalized Markov measure associated with $\mathcal{M}$ and μ,
(iii) M is an equilibrium state for u with $P(u)=0$.
(iv) $U^*F(M)=F(M)$
hold the implications $(i)\Rightarrow(ii)\Rightarrow(iii)\Rightarrow(iv)$.

Proof. $(i)\Rightarrow(ii)$ By Proposition 1 (ii),

$$\sum_{i=1}^{N}\int_{K_i} d(x,x_i)\ d\mu(x) < \infty \text{ for all } x_i\in K_i,\ i=1,...,N,$$

and by Lemma 1 (ii), $M(Y)=1$. Therefore, (ii) follows by Lemma 6 from [17] (in [17], we considered K to be a locally compact space, but this was not used in the proof of Lemma 6).

$(ii)\Rightarrow(iii)$ holds by Lemma 5 from [17] (the same argument as in the proof of Proposition 1 in [17]).

$(iii)\Rightarrow(iv)$ holds by Proposition 2 from [17] (the local compactness was not used in the proof of Proposition 2 from [17]). □

Remark 6 Observe that, by Proposition 3, Theorem 2 is a natural generalization of Theorem 1.

Theorem 3 *Suppose CMS $\mathcal{M}$ has a unique invariant Borel probability measure μ. Then the following hold.*
(i) M is a unique equilibrium state for the energy function u,
(ii) $F(M)=\mu$,
(iii) $h_M(S) = -\sum_{e\in E}\int_{K_{i(e)}} p_e\log p_e\ d\mu$.

Proof. By Lemma 1 (ii), $M(Y) = 1$. Therefore, the claim follows by Theorem 2 form [17] (the local compactness was not used for that). $\Box$

Remark 7 The author would like to point out that a similar entropy formula as that proved in Theorem 3 (iii) plays a central role in the recent book of Wojciech Slomczynski [12].

Now, we intend to show that the images under the coding map of ergodic equilibrium states for u which are absolutely continuous with respect to generalized Markov measures are not only invariant, but also observable.

Definition 8 We call $\mu \in P(K)$ *observable* with $\mathcal{M}$ from $x \in K$ if and only if

$$\frac{1}{n}\sum_{i=1}^{n} \delta_{w_{e_i}\circ\ldots\circ w_{e_1}(x)} \overset{w^*}{\to} \mu \text{ for } P_x\text{-a.e. } \sigma \in \Sigma^+.$$

Note that, by Lebesgues Dominated Convergence Theorem,

$$\frac{1}{n}\sum_{i=1}^{n} U^i f(x) \to \mu(f)$$

for all bounded continuous functions f if μ is observable from x.

First, we need to show that there exist ergodic equilibrium states for u which are absolutely continuous with respect to generalized Markov measures.

In [18], we have proved the following lemma.

Lemma 2 *Let $\Lambda \in ES(u)$ and $\tilde{\Lambda} \in P_S(\Sigma)$ such that $\tilde{\Lambda}$ is absolutely continuous with respect to Λ. Then $\tilde{\Lambda} \in ES(u)$.*

Proof. See [18]. $\Box$

Proposition 4 *Let $\mu \in P(\mathcal{M})$. Then there exists an ergodic equilibrium state Λ for u such that Λ is absolutely continuous with respect to M, where M is a generalized Markov measure associated with $\mathcal{M}$ and μ.*

Proof. By Theorem 2, M is an equilibrium state for u. Since Σ is a compact metric space, it is well known that $P_S(\Sigma)$ is a compact (in the weak* topology) convex space. By the Krein-Milman Theorem, $P_S(\Sigma)$ is a closed convex hull of its extreme points, which are exactly the ergodic measures (see e.g. Theorem 6.10 in [14]). Furthermore, the ergodic measures can be characterized as the minimal elements of the set $P_S(\Sigma)$ with respect to the partial order given by the absolute continuity relation (see e.g. [9], Lemma 2.2.2) (to be precise, the absolute continuity relation is a partial order on the equivalence classes consisting of equivalent measures, but every equivalence class containing an ergodic measure consists of a single element). Therefore, there exists an ergodic $\tilde{\Lambda} \in P_S(\Sigma)$ such that $\tilde{\Lambda}$ is absolutely continuous with respect to M. Thus, by Lemma 2, $\tilde{\Lambda} \in ES(u)$. □

Before we move to the theorem, we need to prove a lemma.

Let $\Sigma^+ := E^{\mathbb{N}}$ endowed with the product topology of the discreet topologies. Let $\nu \in P(K)$. Since $x \longmapsto P_x(Q)$ is Borel measurable for all $Q \in \mathcal{B}(\Sigma^+)$ (see e.g. Lemma 1 in [16]), we can define

$$\tilde{\phi}(\nu)(A \times Q) := \int\limits_A P_x(Q) d\nu(x)$$

for all $A \in \mathcal{B}(K)$ and $Q \in \mathcal{B}(\Sigma^+)$. Then $\tilde{\phi}(\nu)$ extends uniquely to a Borel probability measure on $K \times \Sigma^+$ with

$$\tilde{\phi}(\nu)(\Omega) = \int P_x\left(\left\{\sigma \in \Sigma^+ : (x,\sigma) \in \Omega\right\}\right) d\nu(x)$$

for all $\Omega \in \mathcal{B}(K \times \Sigma^+)$. Note that the set of all $\Omega \subset K \times \Sigma^+$ for which the integrand in the above is measurable forms a Dynkin system which contains all rectangles. Therefore, it is measurable for all $\Omega \in \mathcal{B}(K \times \Sigma^+)$.

Now, consider the following map

$$\begin{array}{rcl} \xi : \Sigma & \longrightarrow & K \times \Sigma^+ \\ \sigma & \longmapsto & (F(\sigma), (\sigma_1, \sigma_2, ...)). \end{array}$$

Lemma 3 *Let $\Lambda \in P_S(\Sigma)$ be an equilibrium state for u such that Λ is absolutely continuous with respect to M where M is a generalized Markov measure associated with $\mathcal{M}$ and $\mu \in P(\mathcal{M})$. Then*

$$\xi(\Lambda) = \tilde{\phi}(F(\Lambda)).$$

Proof. We only need to check that

$$\xi(\Lambda)\left(A \times_1 [e_1, ..., e_n]^+\right) = \tilde{\phi}(F(\Lambda))\left(A \times_1 [e_1, ..., e_n]^+\right)$$

for all cylinder sets ${}_1[e_1, ..., e_n]^+ \subset \Sigma^+$ and $A \in \mathcal{B}(K)$. For such sets

$$\begin{array}{rcl} \xi(\Lambda)\left(A \times_1 [e_1, ..., e_n]^+\right) & = & \Lambda\left(F^{-1}(A) \cap_1 [e_1, ..., e_n]\right) \\ & = & \int\limits_{F^{-1}(A)} 1_{{}_1[e_1,...,e_n]} d\Lambda \end{array}$$

where ${}_1[e_1, ..., e_n] \subset \Sigma$ is the pre-image of ${}_1[e_1, ..., e_n]^+$ under the natural projection. Recall that $F^{-1}(A) \in \mathcal{F}$. Therefore, by Theorem 2,

$$E_M\left(1_{{}_1[e]}|\mathcal{F}\right) = p_e \circ F \ M\text{-a.e. for all } e \in E.$$

Since Λ is absolutely continuous with respect to M,

$$E_\Lambda(1_{{}_1[e]}|\mathcal{F}) = p_e \circ F \ \Lambda\text{-a.e.}, \tag{6}$$

analogously as in the proof of Lemma 7 in [18]. This implies, by the shift invariance of Λ and the pull-out property of the conditional expectation, that

$$E_\Lambda(1_{{}_1[e_1,...,e_n]}|\mathcal{F}) = P_{F(\sigma)}\left({}_1[e_1,...,e_n]^+\right) \ \Lambda\text{-a.e..}$$

Therefore,

$$\begin{aligned} \int\limits_{F^{-1}(A)} 1_{{}_1[e_1,...,e_n]} d\Lambda &= \int\limits_{F^{-1}(A)} P_{F(\sigma)}\left({}_1[e_1,...,e_n]^+\right) d\Lambda(\sigma) \\ &= \int\limits_A P_x\left({}_1[e_1,...,e_n]^+\right) dF(\Lambda)(x), \end{aligned}$$

as desired. $\square$

The following theorem is a little bit more then just the usual strong low of large numbers.

Theorem 4 *Let Λ be an ergodic equilibrium state for u such that Λ is absolutely continuous with respect to M where M is a generalized Markov measure associated with $\mathcal{M}$ and $\mu \in P(\mathcal{M})$. Let $f_e : K \longrightarrow [-\infty, +\infty]$ be Borel measurable such that $f_e|_{K_{i(e)}}$ is bounded and continuous for all $e \in E$. Then, for $F(\Lambda)$-a.e. $x \in K$,*

$$\lim_{n\to\infty} \frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ ... \circ w_{\sigma_1}(x) = \sum_{e\in E} \int\limits_{K_{i(e)}} p_e f_e \ d\mu$$

for P_x-a.e. $\sigma \in \Sigma^+$.

Proof. Since Λ is an equilibrium state for u, $\Lambda(Y) = 1$ (otherwise $h_\Lambda(S)+\Lambda(u) = -\infty < P(u)$). Set

$$v(\sigma) = \begin{cases} \log f_{\sigma_1} \circ F(\sigma) & \text{if } \sigma \in Y \\ -\infty & \text{if } \sigma \in \Sigma \setminus Y. \end{cases} \tag{7}$$

Then, by Birkhoff's Ergodic Theorem,

$$\frac{1}{n} \sum_{k=0}^{n-1} v \circ S^k \to \int v \ d\Lambda \ \Lambda\text{-a.e..}$$

Since $F \circ S^k(\sigma) = w_{\sigma_k} \circ ... \circ w_{\sigma_1}(F(\sigma))$ for all $\sigma \in Y$, we can assume, without loss of generality, that

$$\frac{1}{n} \sum_{k=0}^{n-1} f_{\sigma_{k+1}} \circ w_{\sigma_k} \circ ... \circ w_{\sigma_1}(F(\sigma)) \to \int v \ d\Lambda \text{ for all } \sigma \in Y.$$

By (6),

$$\begin{aligned}\int v\ d\Lambda &= \lim_{n\to\infty}\sum_{e\in E}\int 1_{{}_1[e]}n\wedge f_e\circ F\ d\Lambda = \lim_{n\to\infty}\sum_{e\in E}\int p_e\circ Fn\wedge f_e\circ F\ d\Lambda\\ &= \lim_{n\to\infty}\sum_{e\in E}\int\limits_{K_{i(e)}} p_e n\wedge f_e\ dF(\Lambda) = \sum_{e\in E}\int\limits_{K_{i(e)}} p_e f_e\ dF(\Lambda).\end{aligned}$$

Now, applying the map ξ, we get

$$\frac{1}{n}\sum_{k=0}^{n-1} f_{\sigma_{k+1}}\circ w_{\sigma_k}\circ\ldots\circ w_{\sigma_1}(x)\to\sum_{e\in E}\int\limits_{K_{i(e)}} p_e f_e\ dF(\Lambda)\text{ for all }(x,\sigma)\in\xi(Y).$$

By Lemma 3, $\tilde{\phi}(F(\Lambda))(\xi(Y))=\xi(\Lambda)(\xi(Y))=\Lambda(\xi^{-1}(\xi(Y)))\geq\Lambda(Y)=1$. Since $\tilde{\phi}(F(\Lambda))$ is a probability measure,

$$1 = \tilde{\phi}(F(\Lambda))(\xi(Y)) = \int P_x\left(\left\{\sigma\in\Sigma^+ : (x,\sigma)\in\xi(Y)\right\}\right)dF(\Lambda)(x)$$

Since $0\leq P_x\left(\{\sigma\in\Sigma^+ : (x,\sigma)\in\xi(Y)\}\right)\leq 1$, it is a simple consequence of Lusin's Theorem, that $P_x\left(\{\sigma\in\Sigma^+ : (x,\sigma)\in\xi(Y)\}\right)=1$ for $F(\Lambda)$-a.e. $x\in K$. If we set for such x,

$$Q_x := \left\{\sigma\in\Sigma^+ : (x,\sigma)\in\xi(Y)\right\}.$$

Then $P_x(Q_x)=1$ and

$$\frac{1}{n}\sum_{k=0}^{n-1} f_{\sigma_{k+1}}\circ w_{\sigma_k}\circ\ldots\circ w_{\sigma_1}(x)\to\sum_{e\in E}\int\limits_{K_{i(e)}} p_e f_e\ dF(\Lambda)\text{ for all }\sigma\in Q_x.$$

This completes the proof. □

Corollary 1 *There exists $\mu\in P(K)$ which is observable with $\mathcal{M}$ from μ-a.e. $x\in K$. Moreover, for μ-a.e. $x\in K$,*

$$\lim_{n\to\infty}\frac{1}{n}\log P_x({}_1[\sigma_1,\ldots,\sigma_n]) = \sum_{e\in E}\int\limits_{K_{i(e)}} p_e\log p_e\ d\mu$$

for P_x-a.e. $\sigma\in\Sigma^+$..

Proof. Let f be a bounded continuous function on K. Set $f_e := f$ for all $e\in E$. Then, by Proposition 4 and Theorem 4 and Theorem 2, there exists $\mu\in P(\mathcal{M})$ which is observable from μ-a.e. $x\in K$. Now, set $f_e := \log p_e$ for all $e\in E$. Then, by Theorem 4, for μ-a.e. $x\in K$,

$$\lim_{n\to\infty}\frac{1}{n}\log[p_{\sigma_1}(x)p_{\sigma_2}\circ w_{\sigma_1}(x)\ldots p_{\sigma_n}\circ w_{\sigma_{n-1}}\circ\ldots\circ w_{\sigma_1}(x)] = \sum_{e\in E}\int\limits_{K_{i(e)}} p_e\log p_e\ d\mu$$

for P_x-a.e. $\sigma\in\Sigma^+$. □

Remark 8 Note that Corollary 1 also might have some practical value in case of CMSs on $\mathbb{R}^n$, e.g. which are used for image compression, if μ is also non-singular with respect to the Lebesques measure. The latter happens some times if the images $w_e(K_{i(e)})$ do overlap.

Acknowledgements

I would like to thank Tomasz Szarek for informing me about the reference [4].

References

[1] M. F. Barnsley, S. G. Demko, J. H. Elton and J. S. Geronimo, Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré* **24** (1988) 367-394.

[2] N. Berger, Ch. Hoffman, V. Sidoravicius, Nonuniqueness for specifications in $\ell^{2+\epsilon}$, preprint *math.PR/0312344*.

[3] J. H. Elton, An ergodic theorem for iterated maps, *Ergod. Th. & Dynam. Sys.* **7** (1987) 481-488.

[4] K. Horbacz and T. Szarek, Irreducible Markov systems on polish spaces, *preprint.*

[5] M. Iosifescu, Ş. Grigorescu, *Dependence with complete connections and its applications*, Cambridge Tracts in Mathematics, 96. Cambridge University Press, Cambridge, (1990).

[6] R. Isaac, Markov processes and unique stationary probability measures, *Pacific J. Math.* **12** (1962) 273-286.

[7] A. Johansson and A. Öberg, Square summability of variations of g-functions and uniqueness of g-measures, *Math. Res. Lett.* **10** (2003), 587–601.

[8] M. Keane, Strongly Mixing g-Measures, *Inventiones math.* **16** (1972) 309-324.

[9] G. Keller, Equilibrium states in ergodic theory. London Mathematical Society Student Texts, 42. Cambridge University Press, Cambridge, 1998.

[10] F. Ledrappier, Principe variationnel et systèmes dynamiques symboliques, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **30** (1974), 185-202.

[11] R. D. Mauldin and S. C. Williams, Hausdorff dimension in graph directed constructions, *Tran. AMS* **309** (1988) 811-829.

[12] W. Slomczynski, *Dynamical entropy, Markov operators, and itereated function systems*, Rozprawy Habilitacyjne Uniwersytetu Jagiellońskiego Nr 362, Wydawnictwo Uniwersytetu Jagiellońskiego (2003).

[13] P. Walters, Ruelle's Operator Theorem and g-measures, *Tran. AMS* **214** (1975) 375-387.

[14] P. Walters, *An introduction to Ergodic Theory*, Springer-Verlag, (1982).

[15] I. Werner, Contractive Markov systems, *J. London Math. Soc.* **71** (2005) 236-258.

[16] I. Werner, Coding map for a contractive Markov system, *Math. Proc. Camb. Phil. Soc.* to appear **140** (2) (2006), *preprint math.PR/0504247.*

[17] I. Werner, The generalized Markov measure as an equilibrium state, *Nonlinearity* **18** (2005) 2261-2274.

[18] I. Werner, Contractive Markov systems II, *preprint math.PR/0503633.*